

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

Moderna COVID-19 Vaccine

mRNA-1273 SARS-CoV-2 vaccine

Suspension for intramuscular injection

Multiple Dose Vial, 100 mcg / 0.5mL (per dose)

(contains 10 doses of 0.5 mL)

Active Immunizing Agent

HEALTH CANADA HAS AUTHORIZED THE SALE OF THIS COVID-19 Vaccine UNDER AN INTERIM ORDER

Moderna COVID-19 Vaccine is indicated for:

Active immunization against coronavirus disease 2019 (COVID-19) caused by the SARS-CoV-2 virus in individuals 18 years of age and older.

The use of Moderna COVID-19 Vaccine is permitted under an interim authorization delivered in accordance with section 5 of the COVID-19 Interim order (IO)*. Patients should be advised of the nature of the authorization. The interim authorization is associated with Terms and Conditions that need to be met by the Market Authorization Holder to ascertain the continued quality, safety and efficacy of the product. For further information on authorization under this pathway, please refer to Health Canada's IO Respecting the Importation, Sale and Advertising of Drugs for Use in Relation to COVID-19.

* <https://www.canada.ca/en/health-canada/services/drugs-health-products/covid19-industry/drugs-vaccines-treatments/interim-order-import-sale-advertising-drugs.html#a2.8>

Moderna Therapeutics Inc.
200 Technology Square
Cambridge, MA, USA, 02139

Date of Initial Authorization:
December 23, 2020

Submission Control Number: 244946

TABLE OF CONTENTS

Sections or subsections that are not applicable at the time of authorization are not listed.

TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics.....	4
1.2 Geriatrics.....	4
2 CONTRAINDICATIONS	4
3 Serious warning and precautions	4
4 DOSAGE AND ADMINISTRATION	4
4.1 Dosing Considerations	4
4.2 Recommended Dose and Dosage Adjustment	4
4.3 Reconstitution.....	5
4.4 Administration	5
5 OVERDOSAGE	5
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	6
7 WARNINGS AND PRECAUTIONS	6
7.1 Special Populations	7
7.1.1 Pregnant Women.....	7
7.1.2 Breast-feeding.....	7
7.1.3 Pediatrics.....	7
7.1.4 Geriatrics.....	7
8 ADVERSE REACTIONS	8
8.1 Adverse Reaction Overview	8
8.2 Clinical Trial Adverse Reactions	8
8.4 Post-Market Adverse Reactions.....	14
9 DRUG INTERACTIONS	14
10 CLINICAL PHARMACOLOGY	14
10.1 Mechanism of Action	14
11 STORAGE, STABILITY AND DISPOSAL	14
12 SPECIAL HANDLING INSTRUCTIONS	15

PART II: SCIENTIFIC INFORMATION	16
13 PHARMACEUTICAL INFORMATION	16
14 CLINICAL TRIALS	16
14.1 Study Results.....	18
15 MICROBIOLOGY	18
16 NON-CLINICAL TOXICOLOGY	18
PATIENT MEDICATION INFORMATION	20

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

Moderna COVID-19 Vaccine (mRNA-1273 SARS-CoV-2 vaccine) is indicated for active immunization against coronavirus disease 2019 (COVID-19) caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus in individuals 18 years of age and older.

1.1 Pediatrics

The safety and efficacy of Moderna COVID-19 Vaccine in individuals under 18 years of age has not yet been established. (See [ADVERSE REACTIONS](#), and [CLINICAL TRIALS](#) sections)

1.2 Geriatrics

Clinical studies of Moderna COVID-19 Vaccine include participants 65 years of age and older and their data contributes to the overall assessment of safety and efficacy (See [ADVERSE REACTIONS](#) and [CLINICAL TRIALS](#) sections).

2 CONTRAINDICATIONS

Moderna COVID-19 Vaccine is contraindicated in individuals who are hypersensitive to the active ingredient or to any ingredients in the formulation, including any non-medicinal ingredient, or component of the container. (For a complete listing, see [DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING](#) sections).

3 SERIOUS WARNING AND PRECAUTIONS

At the time of authorization, there are no known serious warnings or precautions associated with this product.

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

Moderna COVID-19 Vaccine is a suspension for intramuscular injection that should be administered by a trained healthcare worker. Moderna COVID-19 Vaccine is a two-dose regimen. The second dose should be administered one month after the first dose.

4.2 Recommended Dose and Dosage Adjustment

Moderna COVID-19 Vaccine should be administered intramuscularly, as two 0.5 mL doses, one month apart.

There are no data available on the interchangeability of Moderna COVID-19 Vaccine with other COVID-19 vaccines to complete the vaccination series. Individuals who have received one dose of Moderna COVID-19 Vaccine should receive a second dose of Moderna COVID-19 Vaccine to complete the vaccination series.

4.3 Reconstitution

Moderna COVID-19 Vaccine must not be reconstituted, mixed with other medicinal products, or diluted.

4.4 Administration

Use aseptic technique for preparation and administration.

Preparation

Thaw each vial before use:

- Thaw in refrigerated conditions between 2°C to 8°C for 2 hours and 30 minutes. Let each vial stand at room temperature for 15 minutes before administering.
- Alternatively, thaw at room temperature between 15°C to 25°C for 1 hour.
- Do not re-freeze vials after thawing.

Swirl the vial gently after thawing and between each withdrawal. Do not shake.

Administration

Moderna COVID-19 Vaccine is a white to off-white suspension. It may contain white or translucent product-related particulates. Inspect Moderna COVID-19 Vaccine vials visually for foreign particulate matter and/or discoloration prior to administration. If either of these conditions exists, the vaccine should not be administered.

Moderna COVID-19 Vaccine should be administered by the intramuscular (IM) route only. Do not inject the vaccine intravascularly, subcutaneously or intradermally. The preferred site is the deltoid muscle of the upper arm. A needle length of ≥ 1 inch should be used as needles < 1 inch may be of insufficient length to penetrate muscle tissue in some adults.

Using aseptic technique, cleanse the vial stopper with a single-use antiseptic swab.

Withdraw each 0.5 mL dose of vaccine from the vial using a new sterile needle and syringe for each injection. The dose in the syringe should be used promptly.

Moderna COVID-19 Vaccine is preservative free. Once a dose is withdrawn from the vial, it should be administered immediately. Once the vial has been entered (needle-punctured), it should be discarded after 6 hours. Do not refreeze. Any unused vaccine or waste material should be disposed of in accordance with local requirements.

5 OVERDOSAGE

In the case of a suspected vaccine overdose, monitoring of vital functions and symptomatic treatment are recommended. Contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1: Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Intramuscular injection	Suspension, (0.20 mg /mL), mRNA, encoding the pre fusion stabilized Spike glycoprotein of 2019 novel Coronavirus (SARS-CoV-2) Multiple dose vial (5 mL, containing 10 doses of 0.5 mL)	<ul style="list-style-type: none">• 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC)• Acetic acid• Cholesterol• Lipid SM-102• PEG2000 DMG 1,2-dimyristoyl-rac-glycerol, methoxy-polyethyleneglycol• Sodium acetate• Sucrose• Tromethamine• Tromethamine hydrochloride• Water for injection

Moderna COVID-19 Vaccine is provided as a white to off-white sterile suspension for intramuscular injection. Moderna COVID-19 Vaccine contains a lipid nanoparticle (LNP) comprised of a messenger ribonucleic acid (mRNA) encoding the pre-fusion stabilized Spike glycoprotein of SARS-CoV-2 virus and four lipids, formulated with the non-medicinal ingredients listed in Table 1. Moderna COVID-19 Vaccine does not contain any preservatives, antibiotics, adjuvants, or human- or animal-derived materials.

Moderna COVID-19 Vaccine is supplied in a multi-dose 10R type I glass vial (each of 5 mL) with a 20 mm Fluro Tec-coated chlorobutyl elastomer stopper, 20 mm flip-off aluminum seal. The vial stopper does not contain natural rubber latex. Vials are packaged in a secondary carton containing a total of ten (10) mRNA-1273 vaccine vials per carton.

To help ensure the traceability of vaccines for patient immunization record-keeping as well as safety monitoring, health professionals should record the time and date of administration, quantity of administered dose (if applicable), anatomical site and route of administration, brand name and generic name of the vaccine, the product lot number and expiry date.

7 WARNINGS AND PRECAUTIONS

The clinical data available for Moderna COVID-19 Vaccine are derived from the COVE Phase 3 study and Phase 1 and Phase 2 studies. Serious and unexpected adverse events may occur that have not been previously reported with Moderna COVID-19 Vaccine use.

As with any vaccine, vaccination with Moderna COVID-19 Vaccine may not protect all recipients.

Individuals may not be optimally protected until after receiving the second dose of the vaccine.

Anaphylaxis

As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of a rare anaphylactic event following the administration of this vaccine.

Acute illness

Consideration should be given to postponing immunization in persons with severe febrile illness or severe acute infection. Persons with moderate or severe acute illness should be vaccinated as soon as the acute illness has improved.

Hematologic-Bleeding

As with other intramuscular injections, Moderna COVID-19 Vaccine should be given with caution in individuals with bleeding disorders, such as haemophilia, or individuals currently on anticoagulant therapy, to avoid the risk of haematoma following the injection, and when the potential benefit clearly outweighs the risk of administration.

Immune

Immunocompromised persons, including individuals receiving immunosuppressant therapy, may have a diminished immune response to the vaccine.

Syncope

Syncope (fainting) can occur following, or even before, any vaccination as a psychogenic response to the needle injection. Procedures should be in place to prevent injury from fainting and manage syncopal reactions.

7.1 Special Populations

7.1.1 Pregnant Women

The safety and efficacy of Moderna COVID-19 Vaccine in pregnant women have not yet been established.

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to Moderna COVID-19 Vaccine during pregnancy. Women who are vaccinated with Moderna COVID-19 Vaccine during pregnancy are encouraged to enroll in the registry by calling 1-866-MODERNA (1-866-663-3762).

7.1.2 Breast-feeding

It is unknown if Moderna COVID-19 Vaccine is excreted in human milk. A risk to the newborns/ infants cannot be excluded. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for immunization against COVID-19.

7.1.3 Pediatrics

The safety and efficacy of Moderna COVID-19 Vaccine in children have not yet been established.

7.1.4 Geriatrics

Clinical studies of Moderna COVID-19 Vaccine include participants 65 years of age and older and their data contributes to the overall assessment of safety and efficacy (See [ADVERSE REACTIONS](#) and [CLINICAL TRIALS](#) sections).

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

The safety profile presented below is based on data generated from an ongoing Phase 3 placebo-controlled clinical study on subjects ≥ 18 years of age.

Solicited adverse reactions were reported more frequently among vaccine subjects than placebo subjects. The most frequently reported adverse reactions after any dose were pain at the injection site (92.0%), fatigue (70.0%), headache (64.7%), myalgia (61.5%) and chills (45.4%). The majority of local and systemic adverse reactions had a median duration of 1 to 3 days.

Overall, there was a higher reported rate of solicited adverse reactions in younger age groups; the incidence of lymphadenopathy (axillary swelling/tenderness), fatigue, headache, myalgia, arthralgia, chills, nausea/vomiting, fever was higher in adults 18 to 64 years of age than in those 65 years of age and above. Solicited adverse reactions were also more frequent after the second dose, compared to the first one, including grade 3 local and systemic adverse reactions (see [Table 2](#), [Table 3](#), [Table 4](#) and [Table 5](#) respectively).

8.2 Clinical Trial Adverse Reactions

Clinical trials are conducted under very specific conditions. The adverse reaction rates observed in the clinical trials; therefore, may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another vaccine. Adverse reaction information from clinical trials may be useful in identifying and approximating rates of adverse vaccine reactions in real-world use.

The safety profile presented below is based on data generated in an ongoing Phase 3, placebo-controlled clinical study on subjects ≥ 18 years of age in which pre-specified cohorts of subjects who were either ≥ 65 years of age or 18 to 64 years of age with comorbid medical conditions were included. At the time of the analysis, the safety analysis set included a total of 30,351 subjects who received at least one dose of Moderna COVID-19 Vaccine (n=15,181) or placebo (n=15,170). Subjects were followed for a median of 92 days from first injection and 63 days from second injection.

Solicited adverse reaction data were collected from Day 1 to Day 7 and reported by participants in an electronic diary (e-Diary) after each dose and on electronic case report forms. Reported solicited local and systemic adverse reactions are presented in [Table 2](#), [Table 3](#), [Table 4](#) and [Table 5](#) respectively.

Table 2: Solicited Local Adverse Reactions Within 7 Days After First and Second Injection by Grade-Participants 18-64 Years of Age (Safety Analysis Set*)

Solicited local AR	Dose 1		Dose 2	
	Vaccine Group n (%) N=11406	Placebo Group n (%) N=11407	Vaccine Group n (%) N=10985	Placebo Group n (%) N=10918
Pain				
Any grade	9908 (86.9)	2177 (19.1)	9873 (89.9)	2040 (18.7)
Grade 3 or 4 ^a	366 (3.2)	23 (0.2)	506 (4.6)	22 (0.2)
Erythema				

Solicited local AR	Dose 1		Dose 2	
	Vaccine Group n (%) N=11406	Placebo Group n (%) N=11407	Vaccine Group n (%) N=10985	Placebo Group n (%) N=10918
Any grade	344 (3.0)	47 (0.4)	982 (8.9)	43 (0.4)
Grade 3 or 4 ^b	34 (0.3)	11 (<0.1)	210 (1.9)	12 (0.1)
Swelling/Induration				
Any grade	767 (6.7)	34 (0.3)	1389 (12.6)	36 (0.3)
Grade 3 or 4 ^b	62 (0.5)	3 (<0.1)	182 (1.7)	4 (<0.1)
Axillary swelling/ Tenderness				
Any grade	1322 (11.6)	567 (5.0)	1775 (16.2)	470 (4.3)
Grade 3 or 4	37 (0.3)	13 (0.1)	46 (0.4)	11 (0.1)

*Safety Analyses Set: all randomized participants who received ≥1 vaccine or control dose.

Note: Adverse reaction data were collected from Day 1 to Day 7 after each dose on the electronic diary (e-Diary) by participants and those collected on the eCRF indicated as solicited adverse reactions.

n= # of participants with specified reaction, percentages are based on n/N

N= number of exposed subjects who submitted any data for the event.

^a Pain - Grade 3: any use of Rx pain reliever/prevents daily activity; Grade 4: requires E.R. visit or hospitalization

^b Erythema and Swelling/Induration - Grade 3: >100mm/>10cm; Grade 4: necrosis/exfoliative dermatitis

^c Axillary Swelling/Tenderness collected as solicited local adverse reaction (i.e., lymphadenopathy: localized axillary swelling or tenderness ipsilateral to the vaccination arm) - Grade 3: any use of Rx pain reliever/prevents daily activity; Grade 4: requires E.R. visit or hospitalization.

Table 3: Solicited Local Adverse Reactions Within 7 Days After First and Second Injection by Grade-Participants 65 Years of Age and Older (Safety Analysis Set*)

Solicited local AR	Dose 1		Dose 2	
	Vaccine Group n (%) N=3762	Placebo Group n (%) N=3748	Vaccine Group n (%) N=3692	Placebo Group n (%) N=3648
Pain				
Any grade	2782 (74.0)	481 (12.8)	3070 (83.2)	437 (12.0)
Grade 3 or 4 ^a	50 (1.3)	32 (0.9)	98 (2.7)	18 (0.5)
Erythema				
Any grade	86 (2.3)	20 (0.5)	275 (7.5)	13 (0.4)
Grade 3 or 4 ^b	8 (0.2)	2 (<0.1)	77 (2.1)	3 (<0.1)
Swelling/Induration				
Any grade	165 (4.4)	18 (0.5)	400 (10.8)	13 (0.4)
Grade 3 or 4 ^b	20 (0.5)	3 (<0.1)	72 (2.0)	7 (0.2)

Solicited local AR	Dose 1		Dose 2	
	Vaccine Group n (%) N=3762	Placebo Group n (%) N=3748	Vaccine Group n (%) N=3692	Placebo Group n (%) N=3648
Axillary swelling/ Tenderness				
Any grade	231 (6.1)	155 (4.1)	315 (8.5)	97 (2.7)
Grade 3 or 4	12 (0.3)	14 (0.4)	21 (0.6)	8 (0.2)

*Safety Analyses Set: all randomized participants who received ≥1 vaccine or control dose.

Note: Adverse reaction data were collected from Day 1 to Day 7 after each dose on the electronic diary (e-Diary) by participants and those collected on the eCRF indicated as solicited adverse reactions.

n= # of participants with specified reaction, percentages are based on n/N

N= number of exposed subjects who submitted any data for the event.

^a Pain - Grade 3: any use of Rx pain reliever/prevents daily activity; Grade 4: requires E.R. visit or hospitalization

^b Erythema and Swelling/Induration - Grade 3: >100mm/>10cm; Grade 4: necrosis/exfoliative dermatitis

^c Axillary Swelling/Tenderness collected as solicited local adverse reaction (i.e., lymphadenopathy: localized axillary swelling or tenderness ipsilateral to the vaccination arm) - Grade 3: any use of Rx pain reliever/prevents daily activity; Grade 4: requires E.R. visit or hospitalization.

Table 4: Solicited Systemic Adverse Reactions Within 7 Days After First and Second Injection by Grade -Participants 18-64 Years of Age (Safety Analysis Set*)

Solicited Systemic AR	Dose 1		Dose 2	
	Vaccine Group n (%) N=11406	Placebo Group n (%) N=11407	Vaccine Group n (%) N=10985	Placebo Group n (%) N=10918
Fatigue				
Any grade	4,384 (38.4)	3,282 (28.8)	7,430 (67.6)	2,687 (24.6)
Grade 3 ^a	120 (1.1)	83 (0.7)	1,174 (10.7)	86 (0.8)
Grade 4 ^b	1 (<0.1)	0 (0)	0 (0)	0 (0)
Headache				
Any grade	4,030 (35.3)	3,304 (29.0)	6,898 (62.8)	2,760 (25.3)
Grade 3 ^c	219 (1.9)	162 (1.4)	553 (5.0)	129 (1.2)
Myalgia				
Any grade	2,699 (23.7)	1,628 (14.3)	6,769 (61.6)	1,411 (12.9)
Grade 3 ^a	73 (0.6)	38 (0.3)	1,113 (10.1)	42 (0.4)
Arthralgia				
Any grade	1,893 (16.6)	1,327 (11.6)	4,993 (45.5)	1,172 (10.7)
Grade 3 ^a	47	29	647	37

Solicited Systemic AR	Dose 1		Dose 2	
	Vaccine Group n (%) N=11406	Placebo Group n (%) N=11407	Vaccine Group n (%) N=10985	Placebo Group n (%) N=10918
	(0.4)	(0.3)	(5.9)	(0.3)
Grade 4 ^b	1 (<0.1)	0 (0)	0 (0)	0 (0)
Chills				
Any grade	1,051 (9.2)	730 (6.4)	5,341 (48.6)	658 (6.0)
Grade 3 ^d	17 (0.1)	8 (<0.1)	164 (1.5)	15 (0.1)
Nausea/vomiting				
Any grade	1,068 (9.4)	908 (8.0)	2,348 (21.4)	801 (7.3)
Grade 3 ^e	6 (<0.1)	8 (<0.1)	10 (<0.1)	8 (<0.1)
Fever				
Any grade	105 (0.9)	37 (0.3)	1,908 (17.4)	39 (0.4)
Grade 3 ^f	10 (<0.1)	1 (<0.1)	184 (1.7)	2 (<0.1)
Grade 4 ^g	4 (<0.1)	4 (<0.1)	12 (0.1)	2 (<0.1)
Use of antipyretic or pain medication	2,656 (23.3)	1,523 (13.4)	6,292 (57.3)	1,248 (11.4)

*Safety Analyses Set: all randomized participants who received ≥ 1 vaccine or control dose.

Note: Adverse reaction data were collected from Day 1 to Day 7 after each dose on the electronic diary (e-Diary) by participants and those collected on the eCRF indicated as solicited adverse reactions.

n= # of participants with specified reaction, percentages are based on n/N

N= number of exposed subjects who submitted any data for the event.

^a Grade 3 fatigue, myalgia, arthralgia: Defined as significant; prevents daily activity.

^b Grade 4 fatigue, arthralgia: Defined as requires emergency room visit or hospitalization.

^c Grade 3 headache: Defined as significant; any use of prescription pain reliever or prevents daily activity.

^d Grade 3 chills: Defined as prevents daily activity and requires medical intervention.

^e Grade 3 nausea/vomiting: Defined as prevents daily activity, requires outpatient intravenous hydration.

^h Grade 3 fever: Defined as $\geq 39.0 - \leq 40.0^\circ\text{C}$ / $\geq 102.1 - \leq 104.0^\circ\text{F}$.

ⁱ Grade 4 fever: Defined as $>40.0^\circ\text{C}$ / $>104.0^\circ\text{F}$.

Table 5: Solicited Systemic Adverse Reactions Within 7 Days After First and Second Injection by Grade -Participants 65 Years of Age and Older (Safety Analysis Set*)

Solicited Systemic AR	Dose 1		Dose 2	
	Vaccine Group n (%) N=3762	Placebo Group n (%) N=3748	Vaccine Group n (%) N=3692	Placebo Group n (%) N=3648
Fatigue				
Any grade	1251 (33.3)	851 (22.7)	2152 (58.3)	716 (19.6)
Grade 3 ^a	30 (0.8)	22 (0.6)	254 (6.9)	20 (0.5)
Headache				
Any grade	921 (24.5)	723 (19.3)	1704 (46.2)	650 (17.8)
Grade 3 ^b	52 (1.4)	34 (0.9)	106 (2.9)	33 (0.9)
Myalgia				
Any grade	742 (19.7)	443 (11.8)	1739 (47.1)	398 (10.9)
Grade 3 ^a	17 (0.5)	9 (0.2)	205 (5.6)	10 (0.3)
Arthralgia				
Any grade	618 (16.4)	456 (12.2)	1291 (35.0)	397 (10.9)
Grade 3 ^a	13 (0.3)	8 (0.2)	123 (3.3)	7 (0.2)
Chills				
Any grade	202 (5.4)	148 (4.0)	1141 (30.9)	151 (4.1)
Grade 3 ^c	7 (0.2)	6 (0.2)	27 (0.7)	2 (<0.1)
Nausea/vomiting				
Any grade	194 (5.2)	166 (4.4)	437 (11.8)	133 (3.6)
Grade 3 ^d	4 (0.1)	4 (0.1)	10 (0.3)	3 (<0.1)
Grade 4 ^e	0 (0)	0 (0)	1 (<0.1)	0 (0)
Fever				
Any grade	10 (0.3)	7 (0.2)	370 (10.0)	4 (0.1)
Grade 3 ^f	1 (<0.1)	1 (<0.1)	18 (0.5)	0 (0)
Grade 4 ^g	0 (0)	2 (<0.1)	1 (<0.1)	1 (<0.1)
Use of antipyretic or pain medication	673 (17.9)	477 (12.7)	1546 (41.9)	329 (9.0)

*Safety Analyses Set: all randomized participants who received ≥ 1 vaccine or control dose.

Note: Adverse reaction data were collected from Day 1 to Day 7 after each dose on the electronic diary (e-Diary) by participants and those collected on the eCRF indicated as solicited adverse reactions.

n= # of participants with specified reaction, percentages are based on n/N

N= number of exposed subjects who submitted any data for the event.

^a Grade 3 fatigue, myalgia, arthralgia: Defined as significant; prevents daily activity.

^b Grade 3 headache: Defined as significant; any use of prescription pain reliever or prevents daily activity.

^c Grade 3 chills: Defined as prevents daily activity and requires medical intervention.

^d Grade 3 Nausea/vomiting: Defined as prevents daily activity, requires outpatient intravenous hydration.

^e Grade 4 Nausea/vomiting: Defined as requires emergency room visit or hospitalization for hypotensive shock.

^f Grade 3 fever: Defined as $\geq 39.0 - \leq 40.0^{\circ}\text{C}$ / $\geq 102.1 - \leq 104.0^{\circ}\text{F}$.

^g Grade 4 fever: Defined as $>40.0^{\circ}\text{C}$ / $>104.0^{\circ}\text{F}$.

Unsolicited Adverse Events

Serious Adverse Events

Serious adverse events were reported in 0.5% of participants who received mRNA-1273 and 0.6% of participants who received a placebo, from the first dose until 28 days following the last vaccination. Serious adverse events were reported in 1% of participants who received mRNA-1273 and 1% of participants who received a placebo, from the first dose until the last observation.

There were no other notable patterns or numerical imbalances between treatment groups for specific categories of adverse events (including other neurologic, neuro-inflammatory, and thrombotic events) that would suggest a causal relationship to Moderna COVID-19 Vaccine.

Three serious adverse events were likely related to the mRNA-1273 vaccine: two cases of facial swelling occurring within 7 days of receiving dose 2, in female patients aged 46 and 51; one case of nausea and vomiting with headaches and fever occurring within 7 days after dose 2 and requiring in-hospital treatment in a 61 y.o. female, with past medical history of headaches with nausea and vomiting requiring hospitalization.

No deaths related to the vaccine were reported in the study.

Non-serious Adverse Events

In the Phase 3 study, unsolicited adverse events occurring within 28 days after each vaccination were reported by 23.9% of subjects who received mRNA-1273, and 21.6% of subjects who received the placebo. These adverse events were predominantly solicited adverse reactions occurring outside of the conventional 7-day monitoring period after the injection (injection site pain, fatigue, headaches, myalgia, etc.). Unsolicited adverse events that occurred in $\geq 1\%$ of study participants who received mRNA-1273 and at a rate at least 1.5-fold higher rate than placebo, were lymphadenopathy related events (1.1% of versus 0.6%). All of the lymphadenopathy events are similar to the axillary swelling/tenderness in the injected arm reported as solicited adverse reactions. Hypersensitivity events were reported in 1.5% of the mRNA-1273 group compared to 1.1% of the placebo group, but this imbalance was mostly due to injection site rash and injection site erythema/swelling occurring more frequently in the mRNA-1273 group. There were no other notable patterns or numerical imbalances between treatment groups for specific categories of non-serious adverse events (including neurologic,

musculoskeletal or inflammatory events) that would suggest a causal relationship to Moderna COVID-19 Vaccine.

8.4 Post-Market Adverse Reactions

There are no post-market adverse drug reactions reported for Moderna COVID-19 Vaccine.

9 DRUG INTERACTIONS

No interaction studies have been performed.

Do not mix Moderna COVID-19 Vaccine with other vaccines/products in the same syringe.

10 CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Moderna COVID-19 Vaccine encodes for the pre-fusion stabilized Spike protein of SARS-CoV-2. After intramuscular injection, cells take up the lipid nanoparticle, effectively delivering the mRNA sequence into cells for expression of the SARS-CoV-2 S antigen. The vaccine induces both neutralizing antibody and cellular immune responses to the spike (S) antigen, which may contribute to protection against COVID-19 disease.

11 STORAGE, STABILITY AND DISPOSAL

Storage Prior to Use

As Displayed on the Vial Labels and Cartons

The Moderna COVID-19 Vaccine multiple-dose vials are stored frozen between -25° to -15°C (-13° to 5°F). Store in the original carton to protect from light.

Additional Storage Information Not Displayed on the Vial Labels and Cartons

Do not store on dry ice or below -40°C (-40°F).

Vials can be stored refrigerated between 2° to 8°C (36° to 46°F) for up to 30 days prior to first use.

Unpunctured vials may be stored between 8° to 25°C (46° to 77°F) for up to 12 hours.

Do not refreeze once thawed.

Thawing Vials Prior To Use

The Moderna COVID-19 Vaccine multiple-dose vial contains a frozen suspension that does not contain a preservative and must be thawed prior to administration.

Remove the required number of vial(s) from storage and thaw each vial before use.

Thaw in refrigerated conditions between 2° to 8°C (36° to 46°F) for 2 hours and 30 minutes. After thawing, let vial stand at room temperature for 15 minutes before administering.

Alternatively, thaw at room temperature between 15° to 25°C (59° to 77°F) for 1 hour.

After thawing, do not refreeze.

Punctured vials

Moderna COVID-19 Vaccine is preservative-free. Once the vial has been entered (needle-punctured), it can be stored at room temperature or refrigerated, but must be discarded after 6 hours. Do not refreeze.

12 SPECIAL HANDLING INSTRUCTIONS

Moderna COVID-19 Vaccine must not be mixed with other medicinal products or diluted. Any unused vaccine or waste material should be disposed of in accordance with local requirements.

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION

Drug Substance

Proper name: mRNA-1273 SARS-CoV-2 vaccine

Chemical name: mRNA-1273 LS (Large Scale) Lipid Nanoparticle (LNP)

Product Characteristics:

Moderna COVID-19 Vaccine is an mRNA-lipid complex [lipid nanoparticle (LNP)] dispersion that contains an mRNA (CX-024414) that encodes for the pre-fusion stabilized Spike glycoprotein of 2019-novel Coronavirus (SARS-CoV-2) and four lipids which act as protectants and carriers of the mRNA. The four lipids are: SM-102 (a custom-manufactured, ionizable lipid); PEG2000-DMG (1,2-dimyristoyl-rac-glycerol, methoxy-polyethyleneglycol); 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC) and cholesterol.

Moderna COVID-19 Vaccine is supplied as a multiple-dose liquid ready-to-use suspension at 0.20 mg/mL for intramuscular administration. Moderna COVID-19 Vaccine is in a 10R clear Type 1 glass vial with a rubber serum stopper and an aluminum seal with flip-off plastic cap. Each vial contains 1.26 mg of CX-024414 mRNA and 24.38 mg of SM-102 LNP as a white to off-white dispersion in preservative-free diluent buffer at pH 7.5. There are 10 doses per vial.

14 CLINICAL TRIALS

14.1 Trial Design and Study Demographics

The safety and efficacy of mRNA-1273 COVID-19 Vaccine were evaluated in a Phase 3 randomized, placebo-controlled, multicentre study in participants 18 years of age and older. A total of 30,351 (15,181 in the mRNA-1273 COVID-19 Vaccine group and N=15,170 in the placebo group) participants were randomized equally to receive 2 doses of mRNA-1273 COVID-19 Vaccine or placebo separated by 28 days. Randomization was stratified by age and risk of severe COVID-19 as follows: ≥ 65 years old, < 65 years old and at increased risk for the complications of COVID-19, and < 65 years old and not at increased risk for the complications of COVID-19.

Pregnant or breastfeeding women and individuals with known history of SARS-CoV-2 infection, immunosuppressive or immunodeficient state, asplenia or recurrent severe infections were excluded from the study. The primary efficacy was symptomatic* COVID-19 infection confirmed by Polymerase Chain Reaction (PCR) and by a clinical adjudication committee. The population for the analysis of the primary efficacy endpoint included participants who did not have evidence of prior infection with SARS-CoV-2 through 14 days after the second dose. Participants are planned to be followed for up to 24 months for assessments of safety and efficacy against COVID-19 disease.

* Symptomatic COVID-19 case definition: At least two of the following systemic symptoms: fever ($\geq 38^{\circ}\text{C}$), chills, myalgia, headache, sore throat, new olfactory and taste disorder(s); or the participant must have experienced at least one of the following respiratory signs/symptoms: cough, shortness of breath or difficulty breathing, or clinical or radiographical evidence of pneumonia; and the participant must have at least one NP swab, nasal swab, or saliva sample (or respiratory sample, if hospitalized) positive for SARS-CoV-2 by RT-PCR. COVID-19 cases were adjudicated by a Clinical Adjudication Committee.

Table 6: Demographic Characteristics – Subjects Without Evidence of Infection Prior to 14 Days After Dose 2 – Evaluable Efficacy Population

	Vaccine Group (N=14134) n (%)	Placebo Group (N= 14073) n (%)	Total (N=28207) n (%)
Sex			
Female	6768 (47.9)	6611 (47.0)	13379 (47.4)
Male	7366 (52.1)	7462 (53.0)	14828 (52.6)
Age (years)			
Mean (SD)	51.6 (15.44)	51.6 (15.54)	51.6 (15.49)
Median	53.0	52.0	53.0
Min, max	18, 95	18, 95	18, 95
Age – Subgroups (years)			
18 to <65	10551 (74.6)	10521 (74.8)	21072 (74.7)
65 and older	3583 (25.4)	3552 (25.2)	7135 (25.3)
Race			
American Indian or Alaska Native	108 (0.8)	111 (0.8)	219 (0.8)
Asian	620 (4.4)	689 (4.9)	1309 (4.6)
Black or African American	1385 (9.8)	1349 (9.6)	2734 (9.7)
Native Hawaiian or Other Pacific Islander	35 (0.2)	31 (0.2)	66 (0.2)
White	11253 (79.6)	11174 (79.4)	22427 (79.5)
Other	299 (2.1)	295 (2.1)	594 (2.1)
Ethnicity			
Hispanic or Latino	2789 (19.7)	2780 (19.8)	5569 (19.7)
Not Hispanic or Latino	11212 (79.3)	11165 (79.3)	22377 (79.3)
Race and Ethnicity			
Non-Hispanic White	9023 (63.8)	8916 (63.4)	17939 (63.6)
Communities of color	5088 (36.0)	5132 (36.5)	10220 (36.2)
Occupational Risk*	11586 (82.0)	11590 (82.4)	23176 (82.2)
Healthcare worker	3593 (25.4)	3581 (25.4)	7174 (25.4)
High Risk Condition**			
One high risk condition present	2616 (18.5)	2591 (18.4)	5207 (18.5)
Two or more high risk conditions present	590 (4.2)	576 (4.1)	1166 (4.1)
No high risk condition	10928 (77.3)	10906 (77.5)	21834 (77.4)
Age and Health Risk for Severe COVID-19***			
18 to <65 years and not at risk	8189 (57.9)	8200 (58.3)	16389 (58.1)
18 to <65 years and at risk	2367 (16.7)	2324 (16.5)	4691 (16.6)
≥ 65 years	3578 (25.3)	3549 (25.2)	7127 (25.3)

* Occupational risk includes: Healthcare Workers; Emergency Response; Retail/Restaurant Operations; Manufacturing and Production; Operations, Warehouse Shipping and Fulfillment centers, Transportation and Delivery Services, Border Protection and Military Personnel Personal care and in-home services; Hospitality and Tourism Workers, Pastoral; Social or Public Health Workers; and Educators and Students.

** High risk for severe COVID-19 is defined as patients who meet at least one of the following criteria (protocol-defined):

- Chronic lung disease (e.g., emphysema and chronic bronchitis, idiopathic pulmonary fibrosis, and cystic fibrosis) or moderate to severe asthma
- Significant cardiac disease (eg, heart failure, coronary artery disease, congenital heart disease, cardiomyopathies, and
- pulmonary hypertension)
- Severe obesity (body mass index ≥ 40 kg/m²)
- Diabetes (Type 1, Type 2 or gestational)
- Liver disease
- Human immunodeficiency virus (HIV) infection

*** Age and health risk for severe COVID-19 is used as stratification factor for randomization.

14.1 Study Results

The analysis of the primary efficacy endpoint included 28,207 participants 18 years of age and older (14,134 in the mRNA-1273 COVID-19 Vaccine group and 14,073 in the placebo group). At the time of the final primary efficacy analysis, participants had been followed for symptomatic COVID 19 disease for a median of 2 months after the second dose, corresponding to 3304.9 person years for the mRNA-1273 COVID-19 Vaccine and 3273.7 person years in the placebo group.

There were 11 confirmed COVID-19 cases identified in the mRNA-1273 COVID-19 Vaccine and 185 in placebo groups, respectively, for the primary efficacy analysis. Compared to placebo, efficacy of mRNA-1273 COVID-19 Vaccine in participants with first COVID-19 occurrence from 14 days after Dose 2 was 94.1% (two-sided 95% confidence interval of 89.3% to 96.8%). In participants 65 years of age and older, efficacy of mRNA-1273 COVID-19 Vaccine was 86.4% (two-sided 95% confidence interval of 61.4% to 95.5%). At the time of primary efficacy analysis, there was a total of 30 severe COVID-19 cases starting 14 days after dose 2, per adjudication committee assessment. All 30 cases were in the placebo group.

15 MICROBIOLOGY

No microbiological information is required for this vaccine product.

16 NON-CLINICAL TOXICOLOGY

General Toxicology: Intramuscular administration of Moderna COVID-19 Vaccine (or other Moderna mRNA investigational vaccines) at doses ranging from 9 to 150 mcg/dose administered once every 2 weeks for up to 6 weeks resulted in transient injection site erythema and edema, body temperature increases, and a generalized systemic inflammatory response. Transient hepatocyte vacuolation and/or Kupffer cell hypertrophy, often observed without liver enzyme elevations, was observed and considered secondary to the systemic inflammatory response. In general, all changes resolved within 2 weeks.

Carcinogenicity: Moderna COVID-19 Vaccine has not been evaluated for carcinogenicity in animals, as carcinogenicity studies were not considered relevant to this vaccine.

Genotoxicity: SM-102, a proprietary lipid component of Moderna COVID-19 Vaccine, is not genotoxic in the bacterial mutagenicity and the human peripheral blood lymphocytes chromosome aberration assays. Two intravenous in vivo micronucleus assays were conducted with mRNA therapies using the same lipid nanoparticle (LNP) formulation as Moderna COVID-19 Vaccine. Equivocal results observed at high systemic concentrations were likely driven by micronuclei formation secondary to elevated body

temperature induced by a LNP-driven systemic inflammatory response. The genotoxic risk to humans is considered to be low due to minimal systemic exposure following intramuscular administration, limited duration of exposure, and the negative in vitro results.

Reproductive and Developmental Toxicology: In a pre- and post-natal developmental toxicity study, 0.2mL of a vaccine formulation containing the same quantity of mRNA (100 µg) and other ingredients included in a single human dose of Moderna COVID-19 Vaccine was administered to female rats by the intramuscular route on four occasions: 28 and 14 days prior to mating, and on gestation days 1 and 13. No vaccine-related adverse effects on female fertility, fetal development or postnatal development were reported in the study.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Moderna COVID-19 Vaccine

mRNA-1273 SARS-CoV-2 vaccine for injection

Health Canada has authorized the sale of this COVID-19 vaccine under an Interim Order.

Read this carefully before you start taking **Moderna COVID-19 Vaccine**. This leaflet is a summary and will not tell you everything about this vaccine. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **Moderna COVID-19 Vaccine**.

What is Moderna COVID-19 Vaccine used for?

Moderna COVID-19 Vaccine is a vaccine used to prevent the coronavirus disease 2019 (COVID-19) caused by the SARS-CoV-2 virus. It can be given to adults aged 18 years and older.

How does Moderna COVID-19 Vaccine work?

Moderna COVID-19 Vaccine works by causing the body to produce its own protection (antibodies) against the SARS-CoV-2 virus that causes the COVID-19 infection. Moderna COVID-19 Vaccine uses a molecule called messenger ribonucleic acid (mRNA) to deliver the set of instructions that cells in the body can use to make antibodies to help fight the virus that causes COVID-19.

The vaccine is given by injection with a needle in the upper arm and will require two doses given one month apart.

As with any vaccine, Moderna COVID-19 Vaccine may not fully protect all those who receive it. Even after you have had both doses of the vaccine, continue to follow the recommendations of local public health officials to prevent spread of COVID-19.

Individuals may not be optimally protected until after receiving the second dose of the vaccine.

You cannot get COVID-19 from this vaccine.

What are the ingredients in Moderna COVID-19 Vaccine?

Medicinal ingredients: mRNA-1273 SARS-CoV-2

Non-medicinal ingredients:

- 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC),
- acetic acid,
- cholesterol,
- PEG2000 DMG (1,2-dimyristoyl-rac-glycerol, methoxy-polyethyleneglycol),
- lipid SM-102,
- sodium acetate,
- sucrose,
- tromethamine
- tromethamine hydrochloride,
- water for injection.

Moderna COVID-19 Vaccine comes in the following dosage forms:

White to off-white suspension for injection, 0.20 mg/mL, provided in a multiple dose vial of 10 doses of 0.5mL, each dose containing 100 micrograms of mRNA.

Do not use Moderna COVID-19 Vaccine if:

- you are allergic to the active substance or any of the other ingredients of this vaccine
 - you have had an allergic reaction to a previous dose of Moderna COVID-19 Vaccine
- you currently have symptoms that could be due to COVID-19. Talk with your healthcare professional about your symptoms and getting a COVID-19 test. Your healthcare professional will advise you when you are able to receive the vaccine

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Moderna COVID-19 Vaccine. Talk about any health conditions or problems you may have, including if you:

- Have any allergies or previous problems following administration of Moderna COVID-19 Vaccine such as an allergic reaction or breathing problems
- Have a bleeding problem, bruise easily or use a blood thinning medication
- Have a high fever or severe infection
- Have any serious illness
- Have a weakened immune system due to a medical condition or are on a medicine that affects your immune system.
- Are pregnant, think you may be pregnant or plan to become pregnant
- Are breastfeeding or plan to breastfeed

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

There is no information on the use of Moderna COVID-19 Vaccine with other vaccines. Tell your healthcare professional if you have recently received any other vaccine.

How is Moderna COVID-19 Vaccine given:

- Your doctor, pharmacist or nurse will inject the vaccine into a muscle (intramuscular injection) in your upper arm
- During and after each injection of the vaccine, your doctor, pharmacist or nurse will watch over you for around 15 minutes to monitor for signs of an allergic reaction.

Usual dose:

Moderna COVID-19 Vaccine will be given to you as two 0.5 mL injections. Each injection will be given on a separate visit 1 month apart. It is very important that you return for the second injection, or the vaccine may not work as well.

Overdose:

In the event of suspected overdose with Moderna COVID-19 Vaccine, contact your regional poison control centre.

Missed Dose:

If you forget to go back to your healthcare professional at the scheduled time for your next dose, ask your healthcare professional for advice.

What are possible side effects from using Moderna COVID-19 Vaccine?

Like all vaccines, Moderna COVID-19 Vaccine can cause side effects

The following are common or very common side effects of Moderna COVID-19 Vaccine. Most of these side effects are mild and do not last long. Tell your doctor if you or your child have side effects that bother you:

- Pain at the injection site
- Tiredness
- Headache
- Muscle ache and stiffness
- Chills
- Fever
- Swelling or redness at the injection site
- Nausea and/or vomiting
- Enlarged lymph nodes

These are not all the possible side effects you may have when taking Moderna COVID-19 Vaccine. If you experience any side effects not listed here, tell your healthcare professional.

Should you develop any serious symptoms or symptoms that could be an allergic reaction, seek medical attention immediately. Symptoms of an allergic reaction include:

- hives (bumps on the skin that are often very itchy)
- swelling of the face, tongue or throat
- difficulty breathing

If you experience a severe allergic reaction, call 9-1-1, or go to the nearest hospital.

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Suspected Side Effects for Vaccines

For the general public: Should you experience a side effect following immunization, please report it to your healthcare professional.

Should you require information related to the management of the side effect, please contact your healthcare professional. The Public Health Agency of Canada, Health Canada and Moderna Therapeutics Inc. cannot provide medical advice.

For healthcare professionals: If a patient experiences a side effect following immunization, please complete the Adverse Events Following Immunization (AEFI) Form appropriate for your

province/territory (<https://www.canada.ca/en/public-health/services/immunization/reporting-adverse-events-following-immunization/form.html>) and send it to your local Health Unit.

Storage:

Do not use this vaccine after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month.

Your doctor or pharmacist is responsible storing, supplying and administering this vaccine, as well as disposing of any unused product correctly.

Keep out of reach and sight of children.

If you want more information about Moderna COVID-19 Vaccine:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website <https://www.modernacovid19global.com/ca/>, or by calling 1-866-MODERNA(1-866-663-3762).

This leaflet was prepared by Moderna Therapeutics Inc.

Last Revised December 23, 2020